

Identify normative values of balance tests toward neurological assessment of sports related concussion

Samaneh Eemanipure, MSc.; Parvaneh Shafinia, Seyed Esmaeel Hashemi-Shaykh Shabani, Ahmad Ghotbi-Varzaneh¹

Shahid Chamran University of Ahvaz, Ahvaz, Iran

Objectives: Deterioration in postural control mechanisms is termed postural instability and results increased postural sway and many laboratory techniques and instruments are characterized by a wide range of neurological signs and symptoms to the medical management. Thus the current study designed to assess the reliability of commonly used clinical measures of balance and determined normal values. Also, the second purpose was scrutiny of effect age, length; weight and body mass index (BMI) on perform clinical balance tests.

Method: One hundred and thirty three participants (18- 59 years), that have at least three time sports activity in one week, performed three timed tests: Time- up and Go (TUG), Tandem Gait (TG), and Walking on Balance Beam (WOBB) on firm surface.

Results: Reliability data were produced for each tests of motor performance. We found that the first performance of three trials was slower, and the relationship between some factors and these battery tests were examined. Means (\pm SD) for each measure were averaged across three trials. Time to complete TG was 13.6 ± 1.1 s. TUG value was 6.9 ± 1.03 and WOBB was 6.9 ± 1.03 s. Conclusions: our results revealed that three clinical balance test batteries - TUG, TG and WOBB tests are the stability measures to assess of sports related concussion. Also, the results of current study appeared that the time to perform these tests was slower than the other studies.

Keywords: normative value, gait, time, reliability

Submitted: 05 Jan 2012

Accepted: 19 Feb 2012

Introduction

Decadence in postural control mechanisms is termed postural instability and results increased postural sway (1). Stabilization of the whole body orientation with respect to the gravity, the support surface, visual surround, muscles and central nervous is a critical component of postural control (2-5). Motor co-ordination (6) and standing balance (3) are essential for many activities of daily living and adequate upper extremity performance. The ability to walk is a quick and inexpensive measure of physical function and an important component of quality of life (7).

Many laboratory techniques that including sports concussion as a common injury in sport (8-10) are characterized by a wide range of neurological signs and symptoms and especially in mild traumatic concussion (11-13). Variety of clinical instruments and symptom checklists used to assess for a sport

concussion with return- to- play decisions to the medical management (14, 15). Several standardized assessment (i.e. standard assessment of concussion (SAC)) or computerized test batteries are commonly used as a measure of impaired cognitive performance post injury. The computer administered test batteries designed for concussion management and return- to- play decisions in athletics are COG sport and immediate post- concussion assessment and cognitive testing (IMPACT). These tasks are described as measuring psycho-motor speed and offered to administration and scoring of test protocols. But these tests have not yet been validated for use in the follow up of sports related concussion (10, 16-18). Recently from combine some validated tools into a single sideline, the sport concussion assessment tool (SCAT) tool were developed by the 1ST and 2ND International Symposiums were held in Vienna, Austria and Prague respectively. This

1. All Correspondence to: Ahmad Ghotbi-Varzaneh; Email: <a.ghotbi@scu.ac.ir>

document is developed for use by health professionals, coaches and other people involved in the care of injured athlete (17). Shuttelworth (2008) proposed that neuro-cognitive evaluation is warranted for any concussive injury to increase diagnostic senility and provide prognostic indications (18).

Some clinical measurement tools have been developed in an attempt to measure dynamic and static balance. These tools were developed to assess balance during a functional performance task in normal participants. Whereas these tests are commonly used as evaluation of neuro-motor function, they have the potential to act as valuable screening items in the assessment of sports related concussion (18). Quantitative gait analysis has been used to illustrate of neurological characteristic features of neurological gait disturbances (19). Number of studies compared gait parameters in wide variety of neurological diseases impair gait and in healthy elderly individuals (19 – 24). Making disequilibrium in patients with few or any neurological signs accounted for gait disturbances (22), and some studies examined the chronic effects of concussion on gait (25). Thus assessment these screening tools are necessary for a sport concussion. In this study, these evaluation tools are Time up and GO (TUG) test (26), Tandem Gait (TG) test and Walking on Balance Beam (WOBB). The last test was new and examined in our study.

Schneiders et al (2010) determined normative values for TG test and demonstrated that TG had excellent reliability in the neurological assessment of sport concussion. Time to complete this test was 11.2 ± 1.2 s in healthy subjects 16-37 years old (27). In another study the reliability coefficients for TG was examined and suggested that further study should be directed toward improving the validity of this test for use with older people (28). In the bischoff et al investigation, it is recorded that 92% of community-dwelling elderly performed the TUG test in less than 12 second. They recommended the TUG test as a screening tool and noted that it is a necessary tool assessment in elderly women (29). Iesles et al stated that normal values for TUG test in subject 20-29 and 30-39 years old were 5.31 ± 0.25 , and 5.39 ± 0.23 , respectively (30).

Therefore, the purpose of this study was to screening of three timed tests of motor performance for used in evaluation of sports concussion and to verify normative values for these tests. The second purpose of this study was to determine the influence of age, sex, leg dominance and body mass index.

Materials and Methods

Convenience samples of individual aged 18-59 years were recruited for this study and one hundred thirty three participants, 50 women ($X=22.1 \pm 1.97$) and 83 men ($X=27.9 \pm 9.45$) were, enrolled. As Schneider's (2010) referrers, according to methods used in povlov et al (2010) and linnert (2000) studies, 120 participants were the enough sample size for calculation of 90% confidence interval and 95% central interval in parametric and non-parametric researches (cited by Schneiders et al., 2010) (27, 31,32). The subjects were chosen from Shahid Chamran university of Ahvaz that had at least three athletic activity sessions in every week. After giving their written informed consent, subjects participated in a structured interview and filled questionnaire and the persons with identified muscle skeletal or neurological disorder, use of drugs that might affect motor tasks and diabetes mellitus were screened from the study. The order effects were randomly identified. 36% of subjects performed TUG test, and 33% executed TG test at first and another one did WOBB test early. Tests performed without shoes and were given rests between repetitions of tests almost 10 seconds and between tests 15 seconds, so that fatigue was not a problem. Also between the second repetitions of the tests in the same day, gave 15 minute rests. In beginning the tests, the subjects were given one untimed trait to insure they understood the tests and then performed three timed trails.

Measurements

The assessment procedure started with measured variables such as height, weight and BMI in the quiet clinic of university. The clinical balance tests used were the TUG test, the TG test and WOBB test that were performed using the protocols described by the original authors excepted of the WOBB test.

In the TUG test, the subjects were seated correctly in a chair with arms, while back of them resting on the back of the chair. There was a marker on the floor 3 meters away from the chair. So that it was easily seen by the subject. On the word GO the subjects standing up from a seated position without using the armchair, walked to the line on the floor, turned around and walked back to the chair and seated down and the stopwatch was stopped. The time interval between starting and stopping the watch provided the time for the test (26).

TG test performed as the same as the time-up-and-go test without the subject seating in a chair. The

starting position was standing with foot together behind a starting line with eyes open, and then the participant walked along a 3 m line in an alternate heel-to-toe fashion, turned and returned to the start point and registered the time of the correct performance.

WOBB test was the new test that examined in the study and used to developed balance and co-ordination. The subjects began the task standing with their feet together behind the Balance Beam and then with preferential foot. Subjects walked along a 5 cm wide, 2.48 m line with short steps and while their hands are free. Subjects failed the test if they deviated from the track. Three trials were recorded.

Data analysis

The testing protocol was performed by two trained and harmonic examiners –examiner 1 and examiner

2. Intra – rater reliability of the motor performance measures are based on the same examiner – examiner 1 – during the same test session and 1 week following the start testing. And inter- rater reliability was assessed by the two examiners both during one performance, simultaneously. We used t tests and general liner model to investigate the influence of several factors -leg dominance, hand dominance, order of testing, age, sex and body max index. The level of significance was set at $P < 0.05$ and $P < 0.01$.

Results

One hundred and thirty three subjects aged 18- 39 years with demographics (50 women, 83 men, weigh: 63.84 ± 10.25 , height: 168.21 ± 5.5 , BMI: 22.55 ± 3.53) participated in the current study. The intra- rater and inter -rater reliability results are summarized in table 1.

Table 1. Intra- class correlation coefficients

Measure	Intrarater-same session(N=133)		Intrarater-between session (N=64)		Inter rater- two examiner(N=57)	
	ICC (single)	ICC (average)	ICC (single)	ICC (average)	ICC (single)	ICC (average)
Time-up-and GO	.764	.895	.764	.855	.992	.996
Tandem Gait	.961	.687	.905	.946	.685	.697
Walking on balance beam	.999	.999	.782	.874	.335	.353

The results revealed absolute reliability for TG test and TUG test and WOBB test that suggesting they

are clinically possible outcome of balance and co-ordination.

Table 2. Individual trial means \pm SD of trials 1-3. Time in seconds (n = 133)

Measure	Trial 1	Trial 2	Trial 3	Mean of 3 Trials
Time- up- and Go	7.2 \pm 1.2	6.9 \pm 1.0	6.8 \pm .9	6.9 \pm 1.03
Tandem Gait	13.5 \pm 1.2	13.8 \pm 1.1	13.5 \pm 1.1	13.6 \pm 1.1
Walking on balance beam	7.2 \pm 1.2	6.9 \pm 1.0	6.8 \pm .9	6.9 \pm 1.03

Also, there was a significant different ($P < 0.001$) between first trial compared to trial 2 and trial 3 in the three test. Also, there was an apparent age and length influence in TG. For Gait Equilibrium test just determined age effect. About TUG also the results shown the significant age, weight and BMI effects were associated with the time to perform TUG. This study noted no order effect for any motor performance measure.

Discussion

In this study TG, WOBB, and TUG proved to be reliable tests and to be used to neurologically screen for sports-related concussion. Using similar protocol, schneiders et al (2010) reported that the

TG was precise and reliable test when administered by the same evaluator. Also, this study supported the results of Schneiders et al (2010) study findings, that the first trial was considerably slower than subsequent trials. But the current study reports values slower for motor performance measured especially in TG. Widespread participant in age range and different between their indexes including age, height, weight, mass and sexes in the two studies might be the elements of these different results.

In The TUG test performance of subjects independent on effects of organ impairments, such as low muscle strength and decreased balance (29) then these tests have enough reliability in different

environments. This clinical measure assess is useful in screening, transition phases associated with balance, such as stand, turn, and sit, as well as gait and can be examined (33). In this test sensory-motor abilities combine and the person with sway in balance have lower constancy in movements. Also there isn't any reported about reliability of WOBB in the past investigation and this study is the first to report normative values for timed versions of this tool.

Non important differences were found in this study between sexes in these tests. In due to significant effect of age in studies (34), it was considered important to utilization age different in performance tests batteries and should be apply in assessments sport concussion. Additionally, in clinical evaluation techniques, repeated trials (35) and learning effects (36) are an important aspect too.

Though sport can play a key role in the lives people, injury incidence is widespread in sport locations. Unfortunately, there are not injury reports in our country -Iran- especially in mild injuries. In one investigation in Taekwondo, as the most popular martial art among Iranian sportsmen, the most frequent injuries were mild (68.8%) and critical injuries (24.7%).and the head and neck also sustained moderate injury (37). Our study is the first tests reliability research in concussion in Iran, and further investigation is necessary to evaluate these

motor performance measures to other health subjects and functional status parameters.

A limitation of our study was the small sample size and left hand and left foot which were investigated in this study. In order to improve the results, further studies with wide and different type's subjects is recommended. Another limitation of this study was that in inter- rater reliability two tester records simultaneously, and the results can influence with this case and these results should be used with caution.

Conclusion

These conclusions provided reliability levels for TG, TUG and WOBB suggesting they are clinically feasible outcome measures of balance in neurological of sports related concussion, and more investigations recommended with using these three tools. Further studies should include a population of subjects with impairment of motor function.

Acknowledgements:

The author wants to thank Azam Robat Sarpushi for giving her the time to help me for performance this work. We also would like to thank all individuals for their participation in this work. This work Supported by Dr Parvaneh Shafinia Fund and Shahid Chamran University Grant.

References

1. Pendergrass TL, Moore JH, Gerber JP. Postural control after a 2-mile run. *Mil Med.* 2003;168(11):896.
2. Deliagina TG, Orlovsky GN, Zelenin PV, Beloozerova IN. Neural bases of postural control. *Physiology.* 2006;21(3):216–25.
3. Horak FB. Postural orientation and equilibrium: what do we need to know about neural control of balance to prevent falls? *Age Ageing.* 2006;35(2):ii7–ii11.
4. Isles RC, Choy NLL, Steer M, Nitz JC. Normal values of balance tests in women aged 20-80. *J Am Geriatr Soc.* 2004;52(8):1367–72.
5. Nishikawa K, Biewener AA, Aerts P, Ahn AN, Chiel HJ, Daley MA, et al. Neuromechanics: an integrative approach for understanding motor control. *Integr Comp Biol.* 2007;47(1):16–54.
6. Desrosiers J, Hebert R, Bravo G, Dutil E. Upper-extremity motor co-ordination of healthy elderly people. *Age Ageing.* 1995;24(2):108–12.
7. Enright PL, Sherrill DL. Reference equations for the six-minute walk in healthy adults. *Am J Respir Crit Care Med.* 1998;158(5 (1)):1384–7.
8. Bailes JE, Hudson V. Classification of Sport-Related Head Trauma: A Spectrum of Mild to Severe Injury. *J Athl Train.* 2001;36(3):236–43.
9. Heilbronner RL, Bush SS, Ravdin LD, Barth JT, Iverson GL, Ruff RM, et al. Neuropsychological consequences of boxing and recommendations to improve safety: a National Academy of Neuropsychology education paper. *Arch Clin Neuropsychol Off J Natl Acad Neuropsychol.* 2009;24(1):11–9.
10. McCrory P, Makkissi M, Davis G, Collie A. Value of neuropsychological testing after head injuries in football. *Br J Sports Med.* 2005;39 (1):i58–63.
11. De Beaumont L, Theoret H, Mongeon D, Messier J, Leclerc S, Tremblay S, et al. Brain function decline in healthy retired athletes who sustained their last sports concussion in early adulthood. *Brain J Neurol.* 2009;132(3):695–708.
12. Blake H, McKinney M, Treece K, Lee E, Lincoln NB. An evaluation of screening measures for cognitive impairment after stroke. *Age Ageing.* 2002;31(6):451–6.
13. Bailes JE, Hudson V. Classification of Sport-Related Head Trauma: A Spectrum of Mild to Severe Injury. *J Athl Train.* 2001;36(3):236–43.
14. Randolph C, Millis S, Barr WB, McCreary M, Guskiewicz KM, Hammeke TA, et al. Concussion symptom inventory: an empirically derived scale for monitoring resolution of symptoms following sport-related concussion. *Arch Clin Neuropsychol Off J Natl Acad Neuropsychol.* 2009;24(3):219–29.
15. Alla S, Sullivan SJ, McCrory P, Schneiders AG, Handcock P. Does exercise evoke neurological symptoms in healthy subjects? *J Sci Med Sport.* 2010;13(1):24–6.
16. Locklin JA. Development of a measure of visumotor control for assessing the long- term effects of concussion. A Thesis in psychology. 2004.
17. Mccrory P, Johnston K, Meeuwisse W, Aubry M, et al. Summary and agreement statement of the 2 nd international

- conference on concussion in sport/ prague 2004. *Clin J Sport Med*. 2005; 15(2):48-55.
18. Shuttleworth-Edwards AB. Central or peripheral?: A positional stance in reaction to the Prague statement on the role of neuropsychological assessment in sports concussion management. *Arch Clin Neuropsychol*. 2008;23(5):479-85.
 19. Ebersbach G, Sojer M, Valldeoriola F, Wissel J, Muller J, Tolosa E, et al. Comparative analysis of gait in Parkinson's disease, cerebellar ataxia and subcortical arteriosclerotic encephalopathy. *Brain J Neurol*. 1999;122 (7):1349-55.
 20. Buchner DM, Cress ME, de Lateur BJ, Esselman PC, Margherita AJ, Price R, et al. The effect of strength and endurance training on gait, balance, fall risk, and health services use in community-living older adults. *J Gerontol A Biol Sci Med Sci*. 1997;52(4):M218-224.
 21. Morris ME, Cantwell C, Vowels L, Dodd K. Changes in gait and fatigue from morning to afternoon in people with multiple sclerosis. *J Neurol Neurosurg Psychiatry*. 2002;72(3):361-5.
 22. Liston R, Mickelborough J, Bene J, Tallis R. A new classification of higher level gait disorders in patients with cerebral multi-infarct states. *Age Ageing*. 2003;32(3):252-8.
 23. Schlicht J, Camaione DN, Owen SV. Effect of intense strength training on standing balance, walking speed, and sit-to-stand performance in older adults. *J Gerontol A Biol Sci Med Sci*. 2001;56(5):M281-286.
 24. Malatesta D, Simar D, Saad HB, Prefaut C, Caillaud C. Effect of an overground walking training on gait performance in healthy 65-to 80-year-olds. *Exp Gerontol*. 2010;45(6):427-34.
 25. Martini DN, Sabin MJ, DePesa SA, Leal EW, Negrete TN, Sosnoff JJ, et al. The chronic effects of concussion on gait. *Arch Phys Med Rehabil*. 2011;92(4):585-9.
 26. Wall JC, Bell C, Campbell S, Davis J. The Timed Get-up-and-Go test revisited: measurement of the component tasks. *J Rehabil Res Dev* 2000;37(1):109-13.
 27. Schneiders AG, Sullivan SJ, Gray AR, Hammond-Tooke GD, McCrory PR. Normative values for three clinical measures of motor performance used in the neurological assessment of sports concussion. *J Sci Med Sport Sports Med Aust*. 2010;13(2):196-201.
 28. Giorgetti MM, Harris BA, Jette A. Reliability of clinical balance outcome measures in the elderly. *Physiother Res Int J Res Clin Phys Ther*. 1998;3(4):274-83.
 29. Bischoff HA, Stahelin HB, Monsch AU, Iversen MD, Weyh A, von Dechend M, et al. Identifying a cut-off point for normal mobility: a comparison of the timed "up and go" test in community-dwelling and institutionalised elderly women. *Age Ageing*. 2003;32(3):315-20.
 30. Isles RC, Choy NLL, Steer M, Nitz JC. Normal values of balance tests in women aged 20-80. *J Am Geriatr Soc*. 2004;52(8):1367-72.
 31. Pavlov IY, Wilson AR, Delgado JC. Resampling approach for determination of the method for reference interval calculation in clinical laboratory practice. *Clin Vaccine Immunol Cvi*. 2010;17(8):1217-22.
 32. Linnert K. Nonparametric estimation of reference intervals by simple and bootstrap-based procedures. *Clin Chem*. 2000;46(6(1)):867-9.
 33. Newton RA. Balance screening of an inner city older adult population. *Arch Phys Med Rehabil*. 1997;78(6):587-91.
 34. Stevens KN, Lang IA, Guralnik JM, Melzer D. Epidemiology of balance and dizziness in a national population: findings from the English Longitudinal Study of Ageing. *Age Ageing*. 2008 ;37(3):300-5.
 35. Riemann BL, Guskiewicz KM. Effects of mild head injury on postural stability as measured through clinical balance testing. *J Athl Train*. 2000;35(1):19-25.
 36. Schneiders AG, Sullivan SJ, McCrory PR, Gray A, Maruthayanar S, Singh P, et al. The effect of exercise on motor performance tasks used in the neurological assessment of sports-related concussion. *Br J Sports Med*. 2008;42(12):1011-3.
 37. Ziaee V, Rahmani SH, Rostami M. Injury rates in Iranian taekwondo athletes; a prospective study. *Asian J Sports Med*. 2010;1(1):23-8.