Letter to Editor





The Imperative Role of Facial Rehabilitation Surgeries in Stroke Patients

Hashem Shemshadi1* 6

1. Department of Clinical Sciences, School of Behavioral Sciences, University of Social Welfare and Rehabilitation Sciences, Tehran, Iran.



cardiovascular accidents.

Citation Shemshadi H. The Imperative Role of Facial Rehabilitation Surgeries in Stroke Patients. Iranian Rehabilitation Journal. 2025; 23(3):245-246. http://dx.doi.org/10.32598/irj.23.3.254.6



am writing this letter to highlight
the profound significance of facial
rehabilitation surgeries for patients who
have endured the debilitating effects
of stroke. This practice, deeply rooted
in evidence-based clinical rehabilitation, serves as an
alternative modality for patient intervention following

Stroke, a leading cause of long-term disability, often leaves survivors grappling with myriad physical and emotional challenges. Among these, facial neuro-anatomical changes are common and significantly impact the quality of life (QoL) of affected individuals. In this context, facial rehabilitation surgeries have emerged as a beacon of hope, offering a path towards recovery and improved well-being.

Patients who have navigated two years post their cardiovascular accidents are deemed suitable candidates for this rehabilitation procedure. These individuals undergo rigorous clinical and para-clinical evaluations to ensure their stability and readiness for reanimation surgery. It is crucial that the underlying causative diseases for the illness above, along with any other potential systemic ailments, are under comprehensive medical control.

Stroke patients, often grappling with facial neuroanatomical changes, are frequently cherished by relatives and friends. However, the natural sense of loss experienced by these patients can trigger a spectrum of emotional disorders. This can manifest as poor communication, impaired speech, and strained interactions with family, friends, and the surrounding environment. A pervasive feeling of hopelessness is not uncommon among these patients, which can further impede their willingness and ability to perform basic self-care.

Facial rehabilitation reanimation surgery is a valuable method to remodel the face, particularly when most neuromuscular groups are hindered by stroke. This surgical intervention can significantly enhance the patient's QoL, restore a sense of normalcy, and boost their self-esteem.

I urge my esteemed colleagues to turn their attention to this imperative subject, underscore the importance of this idea, and initiate further research into the issues above. This subject may also serve as a focal point for developing additional rehabilitation alternatives, creating educational course plans for rehabilitation students, increasing social awareness, and benefiting patients. Elderly patients with this disease may particularly benefit from this rehabilitation surgical modality, as it can significantly enhance their QoL.

* Corresponding Author:

Hashem Shemshadi, Professor.

Address: Department of Clinical Sciences, School of Behavioral Sciences, University of Social Welfare and Rehabilitation Sciences, Tehran, Iran. Tel: +98 (21) 22180140

E-mail: shemshadii@gmail.com



Copyright © 2025 The Author(s)

In conclusion, facial rehabilitation surgery is of paramount importance in stroke patients. This field is ripe for exploration and innovation, with the potential to transform countless lives. As we continue to advance our understanding and capabilities, it is our collective responsibility to ensure that these advancements translate into improved patient care and outcomes.